

Excited light isoscalar mesons from lattice QCD

Christopher Thomas, Jefferson Lab

thomasc@jlab.org

Lattice 2011, Squaw Valley, CA, July 2011

With Jo Dudek, Robert Edwards, Mike Peardon, Bálint Joó,
David Richards and the *Hadron Spectrum Collaboration*



Outline

- Introduction
- Method outline
- Results – isoscalar meson spectrum
- Summary and outlook

PR D83, 111502 (2011)

Motivation

Upcoming experimental efforts in light meson sector (and charmonium)

GlueX and CLAS12 (JLab), BESIII, PANDA, ...

Motivation

Upcoming experimental efforts in light meson sector (and charmonium)


GlueX and CLAS12 (JLab), BESIII, PANDA, ...

Quark-antiquark pair: $2S+1L_J$

Parity: $P = (-1)^{L+1}$

Charge Conj Sym: $C = (-1)^{L+S}$

$J^{PC} = 0^{-+}, 0^{++}, 1^{--}, 1^{++}, 1^{+-}, 2^{--}, 2^{++}, 2^{-+}, \dots$



Motivation

Upcoming experimental efforts in light meson sector (and charmonium)


GlueX and CLAS12 (JLab), BESIII, PANDA, ...

Quark-antiquark pair: $2S+1L_J$

Parity: $P = (-1)^{(L+1)}$

Charge Conj Sym: $C = (-1)^{(L+S)}$

$J^{PC} = 0^{-+}, 0^{++}, 1^{--}, 1^{++}, 1^{+-}, 2^{--}, 2^{++}, 2^{-+}, \dots$



Exotics ($J^{PC} = \mathbf{1}^{-+}, \mathbf{2}^{+-}, \dots$)? – can't just be a $q\bar{q}$ pair


Probe low energy d.o.f. of QCD

e.g. hybrids, multi-mesons

Motivation

Upcoming experimental efforts in light meson sector (and charmonium)

GlueX and CLAS12 (JLab), BESIII, PANDA, ...




Exotics ($J^{PC} = \mathbf{1}^{-+}, \mathbf{2}^{+-}, \dots$)? – can't just be a $q\bar{q}$ pair

Probe low energy d.o.f. of QCD

e.g. hybrids, multi-mesons

Motivation

Up



Exotics ($J^{PC} = \mathbf{1}^{-+}, \mathbf{2}^{+-}, \dots$)? – can't just be a $q\bar{q}$ pair

Probe low energy d.o.f. of QCD

e.g. hybrids, multi-mesons

Photoproduction at GlueX/CLAS12

– systematic study of light mesons, particular interest in exotics

Isoscalars

Isoscalars ($I = 0$) e.g. η , η' , ω , ϕ

Isoscalars

Isoscalars ($I = 0$) e.g. η , η' , ω , ϕ

Probe annihilation dynamics in QCD

Isoscalars

Isoscalars ($I = 0$) e.g. η , η' , ω , ϕ

Probe annihilation dynamics in QCD

$m_s = m_u = m_d$ [SU(3) sym]
– eigenstates are octet, singlet

$$\frac{1}{\sqrt{6}} (u\bar{u} + d\bar{d} - 2s\bar{s})$$

$$\frac{1}{\sqrt{3}} (u\bar{u} + d\bar{d} + s\bar{s})$$

Isoscalars

Isoscalars ($I = 0$) e.g. η , η' , ω , ϕ

Probe annihilation dynamics in QCD

$m_s = m_u = m_d$ [SU(3) sym]
– eigenstates are octet, singlet

$$\frac{1}{\sqrt{6}} (u\bar{u} + d\bar{d} - 2s\bar{s})$$

$$\frac{1}{\sqrt{3}} (u\bar{u} + d\bar{d} + s\bar{s})$$

$m_s \neq m_u = m_d$ – physical states are a mixture

‘Ideal mixing’

$$\ell\bar{\ell} \equiv \frac{1}{\sqrt{2}} (u\bar{u} + d\bar{d})$$

$s\bar{s}$

Isoscalars

Isoscalars ($I = 0$) e.g. η , η' , ω , ϕ

Probe annihilation dynamics in QCD

$m_s = m_u = m_d$ [SU(3) sym]
– eigenstates are octet, singlet

$$\frac{1}{\sqrt{6}} (u\bar{u} + d\bar{d} - 2s\bar{s})$$

$$\frac{1}{\sqrt{3}} (u\bar{u} + d\bar{d} + s\bar{s})$$

$$\alpha = 54.7^\circ$$

$m_s \neq m_u = m_d$ – physical states are a mixture

'Ideal mixing'

$$l\bar{l} \equiv \frac{1}{\sqrt{2}} (u\bar{u} + d\bar{d})$$

$s\bar{s}$

$$\alpha = 0$$

In general

$$|a\rangle = \cos \alpha |l\bar{l}\rangle - \sin \alpha |s\bar{s}\rangle$$

$$|b\rangle = \sin \alpha |l\bar{l}\rangle + \cos \alpha |s\bar{s}\rangle$$

Isoscalars

Isoscalars ($I = 0$) e.g. η , η' , ω , ϕ

Probe annihilation dynamics in QCD

$m_s = m_u = m_d$ [SU(3) sym]
– eigenstates are octet, singlet

$$\frac{1}{\sqrt{6}} (u\bar{u} + d\bar{d} - 2s\bar{s})$$

$$\frac{1}{\sqrt{3}} (u\bar{u} + d\bar{d} + s\bar{s})$$

$$\alpha = 54.7^\circ$$

$m_s \neq m_u = m_d$ – physical states are a mixture

‘Ideal mixing’

$$|\ell\bar{\ell}\rangle \equiv \frac{1}{\sqrt{2}} (u\bar{u} + d\bar{d})$$

$|s\bar{s}\rangle$

$$\alpha = 0$$

In general

$$|a\rangle = \cos \alpha |\ell\bar{\ell}\rangle - \sin \alpha |s\bar{s}\rangle$$

$$|b\rangle = \sin \alpha |\ell\bar{\ell}\rangle + \cos \alpha |s\bar{s}\rangle$$

Experimentally

ω , ϕ (1^{--}) and $f_2(1270)$, $f_2'(1525)$ (2^{++}) – close to ‘ideal’

η , η' (0^{-+}) – closer to octet-singlet

Isoscalars

Isoscalars ($I = 0$) e.g. η , η' , ω , ϕ

Probe annihilation dynamics in QCD

$m_s = m_u = m_d$ [SU(3) sym]
– eigenstates are octet, singlet

$$\frac{1}{\sqrt{6}} (u\bar{u} + d\bar{d} - 2s\bar{s})$$

$$\frac{1}{\sqrt{3}} (u\bar{u} + d\bar{d} + s\bar{s})$$

$$\alpha = 54.7^\circ$$

$m_s \neq m_u = m_d$ – physical states are a mixture

‘Ideal mixing’

$$|\ell\bar{\ell}\rangle \equiv \frac{1}{\sqrt{2}} (u\bar{u} + d\bar{d})$$

$|s\bar{s}\rangle$

$$\alpha = 0$$

In general

$$|a\rangle = \cos \alpha |\ell\bar{\ell}\rangle - \sin \alpha |s\bar{s}\rangle$$

$$|b\rangle = \sin \alpha |\ell\bar{\ell}\rangle + \cos \alpha |s\bar{s}\rangle$$

Can also mix
with glueballs

Experimentally

ω , ϕ (1^{--}) and $f_2(1270)$, $f_2'(1525)$ (2^{++}) – close to ‘ideal’

η , η' (0^{-+}) – closer to octet-singlet

Variational Method

$$C_{ij}(t) = \langle 0 | O_i(t) O_j(0) | 0 \rangle$$

definite J^{PC}

$$O(t) = \sum_{\vec{x}} e^{i\vec{p}\cdot\vec{x}} \bar{\psi}(x) \Gamma_i \overleftrightarrow{D}_j \overleftrightarrow{D}_k \dots \psi(x)$$

Here up to 3 derivs
with $\mathbf{p} = \mathbf{0}$

Variational Method

$$C_{ij}(t) = \langle 0 | O_i(t) O_j(0) | 0 \rangle$$

definite J^{PC}

$$O(t) = \sum_{\vec{x}} e^{i\vec{p}\cdot\vec{x}} \bar{\psi}(x) \Gamma_i \overleftrightarrow{D}_j \overleftrightarrow{D}_k \dots \psi(x)$$

Here up to 3 derivs
with $\mathbf{p} = \mathbf{0}$

Large basis of operators \rightarrow matrix of correlators

$$C_{ij}(t) v_j^{(n)} = \lambda^{(n)}(t) C_{ij}(t_0) v_j^{(n)}$$

generalised
eigenvector problem

Variational Method

$$C_{ij}(t) = \langle 0 | O_i(t) O_j(0) | 0 \rangle$$

definite J^{PC}

$$O(t) = \sum_{\vec{x}} e^{i\vec{p}\cdot\vec{x}} \bar{\psi}(x) \Gamma_i \overleftrightarrow{D}_j \overleftrightarrow{D}_k \dots \psi(x)$$

Here up to 3 derivs
with $\mathbf{p} = \mathbf{0}$

Large basis of operators \rightarrow matrix of correlators

$$C_{ij}(t) v_j^{(n)} = \lambda^{(n)}(t) C_{ij}(t_0) v_j^{(n)}$$

generalised
eigenvector problem

Eigenvalues \rightarrow energies

$$\lambda^{(n)}(t) \rightarrow e^{-E_n(t-t_0)}$$

$(t \gg t_0)$

$Z^{(n)}$ related to eigenvectors

$$Z_i^{(n)} \equiv \langle 0 | O_i | n \rangle$$

$$\Omega^{(n)} \sim \sum_i v_i^{(n)} O_i$$

Variational Method

$$C_{ij}(t) = \langle 0 | O_i(t) O_j(0) | 0 \rangle$$

definite J^{PC}

$$O(t) = \sum_{\vec{x}} e^{i\vec{p}\cdot\vec{x}} \bar{\psi}(x) \Gamma_i \overleftrightarrow{D}_j \overleftrightarrow{D}_k \dots \psi(x)$$

Here up to 3 derivs
with $\mathbf{p} = \mathbf{0}$

Large basis of operators \rightarrow matrix of correlators

$$C_{ij}(t) v_j^{(n)} = \lambda^{(n)}(t) C_{ij}(t_0) v_j^{(n)}$$

generalised
eigenvector problem

Eigenvalues \rightarrow energies

$$\lambda^{(n)}(t) \rightarrow e^{-E_n(t-t_0)}$$

$(t \gg t_0)$

$Z^{(n)}$ related to eigenvectors

$$Z_i^{(n)} \equiv \langle 0 | O_i | n \rangle$$

$$\Omega^{(n)} \sim \sum_i v_i^{(n)} O_i$$

Var. method uses orthog of eigenvectors; don't just rely on separating energies

Isoscalars in LQCD

Use variational method with large basis of operators

Isoscalars in LQCD

Use variational method with large basis of operators

Basis doubled in size c.f. isovectors:

No glueball ops for now

$$O^l \sim \frac{1}{\sqrt{2}} (\bar{u}\Gamma u + \bar{d}\Gamma d)$$

$$O^s \sim \bar{s}\Gamma s$$

Isoscalars in LQCD

Use variational method with large basis of operators



Basis doubled in size c.f. isovectors:

No glueball ops for now

$$O^l \sim \frac{1}{\sqrt{2}} (\bar{u}\Gamma u + \bar{d}\Gamma d)$$

$$O^s \sim \bar{s}\Gamma s$$

Disconnected diagrams



$$C_{AB}^{q'q}(t', t) = \langle 0 | \mathcal{O}_A^{q'}(t') \mathcal{O}_B^{q\dagger}(t) | 0 \rangle$$

$$C = \begin{pmatrix} -C^{ll} + 2\mathcal{D}^{ll} & \sqrt{2}\mathcal{D}^{ls} \\ \sqrt{2}\mathcal{D}^{sl} & -C^{ss} + \mathcal{D}^{ss} \end{pmatrix}$$

Isoscalars in LQCD

Use variational method with large basis of operators


Basis doubled in size c.f. isovectors:

No glueball ops for now

$$O^l \sim \frac{1}{\sqrt{2}} (\bar{u}\Gamma u + \bar{d}\Gamma d)$$

$$O^s \sim \bar{s}\Gamma s$$

Disconnected diagrams



$$C_{AB}^{q'q}(t', t) = \langle 0 | \mathcal{O}_A^{q'}(t') \mathcal{O}_B^{q\dagger}(t) | 0 \rangle$$

$$C = \begin{pmatrix} -C^{ll} + 2\mathcal{D}^{ll} & \sqrt{2}\mathcal{D}^{ls} \\ \sqrt{2}\mathcal{D}^{sl} & -C^{ss} + \mathcal{D}^{ss} \end{pmatrix}$$

Isoscalars in LQCD

Use variational method with large basis of operators


Basis doubled in size c.f. isovectors:

No glueball ops for now

$$O^l \sim \frac{1}{\sqrt{2}} (\bar{u}\Gamma u + \bar{d}\Gamma d)$$

$$O^s \sim \bar{s}\Gamma s$$

Disconnected diagrams



$$C_{AB}^{q'q}(t', t) = \langle 0 | \mathcal{O}_A^{q'}(t') \mathcal{O}_B^{q\dagger}(t) | 0 \rangle$$

$$C = \begin{pmatrix} -C^{ll} + 2\mathcal{D}^{ll} & \sqrt{2}\mathcal{D}^{ls} \\ \sqrt{2}\mathcal{D}^{sl} & -C^{ss} + \mathcal{D}^{ss} \end{pmatrix}$$

Difficulties with traditional methods

Distillation

PR D80, 054506

Distillation

$$\square_{xy}(t) = \sum_{k=1}^N v_x^{(k)}(t) v_y^{(k)\dagger}(t)$$

$$\square(t) = V(t)V^\dagger(t)$$

$$C_{AB}^{q'q}(t', t) = \langle 0 | \mathcal{O}_A^{q'}(t') \mathcal{O}_B^{q\dagger}(t) | 0 \rangle$$

Distillation

PR D80, 054506

Distillation

$$\square_{xy}(t) = \sum_{k=1}^N v_x^{(k)}(t) v_y^{(k)\dagger}(t)$$

$$\square(t) = V(t) V^\dagger(t)$$

$$C_{AB}^{q'q}(t', t) = \langle 0 | \mathcal{O}_A^{q'}(t') \mathcal{O}_B^{q\dagger}(t) | 0 \rangle$$

$$C_{AB}^{q'q}(t', t) = \langle 0 | \bar{q}' \square(t') \Gamma_{t'}^A \square(t') q'(t') \cdot \bar{q} \square(t) \Gamma_t^B \square(t) q(t) | 0 \rangle$$

Distillation

PR D80, 054506

Distillation

$$\square_{xy}(t) = \sum_{k=1}^N v_x^{(k)}(t) v_y^{(k)\dagger}(t)$$

$$\square(t) = V(t) V^\dagger(t)$$

$$C_{AB}^{q'q}(t', t) = \langle 0 | \mathcal{O}_A^{q'}(t') \mathcal{O}_B^{q\dagger}(t) | 0 \rangle$$

$$C_{AB}^{q'q}(t', t) = \langle 0 | \bar{q}' \square(t') \Gamma_{t'}^A \square(t') q'(t') \cdot \bar{q} \square(t) \Gamma_t^B \square(t) q(t) | 0 \rangle$$

$$C = \begin{pmatrix} -c^{ll} + 2\mathcal{D}^{ll} & \sqrt{2}\mathcal{D}^{ls} \\ \sqrt{2}\mathcal{D}^{sl} & -c^{ss} + \mathcal{D}^{ss} \end{pmatrix}$$

Distillation

PR D80, 054506

Distillation

$$\square_{xy}(t) = \sum_{k=1}^N v_x^{(k)}(t) v_y^{(k)\dagger}(t)$$

$$\square(t) = V(t) V^\dagger(t)$$

$$C_{AB}^{q'q}(t', t) = \langle 0 | \mathcal{O}_A^{q'}(t') \mathcal{O}_B^{q\dagger}(t) | 0 \rangle$$

$$C_{AB}^{q'q}(t', t) = \langle 0 | \bar{q}' \square(t') \Gamma_{t'}^A \square(t') q'(t') \cdot \bar{q} \square(t) \Gamma_t^B \square(t) q(t) | 0 \rangle$$



$$C = \begin{pmatrix} -C^{ll} + 2\mathcal{D}^{ll} & \sqrt{2}\mathcal{D}^{ls} \\ \sqrt{2}\mathcal{D}^{sl} & -C^{ss} + \mathcal{D}^{ss} \end{pmatrix}$$

$$C_{AB}^{q'q}(t', t) = \delta_{qq'} \text{Tr}[\Phi^A(t') \tau_{q'}(t', t) \Phi^B(t) \tau_q(t, t')]$$

$$\mathcal{D}_{AB}^{q'q}(t', t) = \text{Tr}[\Phi^A(t') \tau_{q'}(t', t')] \text{Tr}[\Phi^B(t) \tau_q(t, t)]$$

$$\tau_q(t', t) = V_t^\dagger M_q^{-1}(t', t) V_t$$

$$\Phi^A(t) = V_t^\dagger \Gamma_t^A V_t$$



Isoscalars in LQCD

Use distillation

PR D80, 054506

Number of inversions $\sim N_{\text{vecs}}$ instead of $(N_{\text{colours}} \times L_s^3)$

Isoscalars in LQCD

Use distillation

PR D80, 054506

Number of inversions $\sim N_{\text{vecs}}$ instead of $(N_{\text{colours}} \times L_s^3)$

Use GPUs

Isoscalars in LQCD

Use distillation

PR D80, 054506

Number of inversions $\sim N_{\text{vecs}}$ instead of $(N_{\text{colours}} \times L_s^3)$

Use GPUs

Efficient computations with **large basis of operators**

Isoscalars in LQCD

Use distillation

PR D80, 054506

Number of inversions $\sim N_{\text{vecs}}$ instead of $(N_{\text{colours}} \times L_s^3)$

Use GPUs

Efficient computations with **large basis of operators**

Project onto **definite mom.** at source and sink

Isoscalars in LQCD

Use distillation

PR D80, 054506

Number of inversions $\sim N_{\text{vecs}}$ instead of $(N_{\text{colours}} \times L_s^3)$

Use GPUs

Efficient computations with **large basis of operators**

Project onto **definite mom.** at source and sink

Increase stats. by averaging

$N_t=32, L_t=128$

$$C^{qq}(t) = \frac{1}{N_t} \sum_{t_0}^{N_t} C^{qq}(t + t_0, t_0)$$

$$\mathcal{D}^{q'q}(t) = \frac{1}{L_t} \sum_{t_0}^{L_t} \mathcal{D}^{q'q}(t + t_0, t_0)$$

Isoscalars in LQCD

Use distillation

PR D80, 054506

Number of inversions $\sim N_{\text{vecs}}$ instead of $(N_{\text{colours}} \times L_s^3)$

Use GPUs

Efficient computations with **large basis of operators**

Project onto **definite mom.** at source and sink

Increase stats. by averaging

$N_t=32, L_t=128$

$$C^{qq}(t) = \frac{1}{N_t} \sum_{t_0}^{N_t} C^{qq}(t + t_0, t_0)$$

$$\mathcal{D}^{q'q}(t) = \frac{1}{L_t} \sum_{t_0}^{L_t} \mathcal{D}^{q'q}(t + t_0, t_0)$$

Anisotropic Clover ($a_s/a_t = 3.5$), $a_s \sim 0.12$ fm; 16^3 (2.0 fm)

Lattice details in:
PR D78 054501,
PR D79 034502

$N_f = 2+1, M_\pi \approx 400$ MeV


$N_{\text{vecs}} = 64$

479 cfigs

Correlators


PR D83, 111502 (2011)

$$\mathcal{O}_A = \mathcal{O}_B \sim \gamma_5$$




Isoscalars

PR D83, 111502 (2011)



Isoscalars

PR D83, 111502 (2011)



Flavour Mixings

Determine mixing of physical states by looking at overlaps

$$O^{\ell} \sim \frac{1}{\sqrt{2}} (\bar{u}\Gamma u + \bar{d}\Gamma d)$$

$$O^s \sim \bar{s}\Gamma s$$

$$Z_{A,\ell}^{(n)} = \langle 0 | O_A^{\ell} | n \rangle$$

$$Z_{A,s}^{(n)} = \langle 0 | O_A^s | n \rangle$$

Flavour Mixings

Determine mixing of physical states by looking at overlaps

$$O^{\ell} \sim \frac{1}{\sqrt{2}} (\bar{u}\Gamma u + \bar{d}\Gamma d)$$

$$O^s \sim \bar{s}\Gamma s$$


$$Z_{A,\ell}^{(n)} = \langle 0 | O_A^{\ell} | n \rangle$$

$$Z_{A,s}^{(n)} = \langle 0 | O_A^s | n \rangle$$

$$\alpha_A = \tan^{-1} \left[\sqrt{\frac{Z_{\ell,A}^{(b)} Z_{s,A}^{(a)}}{Z_{\ell,A}^{(a)} Z_{s,A}^{(b)}}} \right]$$

Isoscalars

PR D83, 111502 (2011)




First light isoscalar exotics in LQCD

Most close to ideally mixed

Isoscalars

PR D83, 111502 (2011)




First light isoscalar exotics in LQCD

Most close to ideally mixed

But $\eta - \eta' = 42(1)^\circ$, $f_1 - f_1' = 31(2)^\circ$, 1^{+-} exotics

Isoscalars

PR D83, 111502 (2011)



Most close to ideally mixed

But $\eta - \eta' = 42(1)^{\circ}$, $f_1 - f_1' = 31(2)^{\circ}$, 1^{-+} exotics

Summary and Outlook

Summary

- **Extensive** isoscalar meson spectrum
- **Exotics** and non-exotic **hybrids**, high spin, excited states
- Carefully constructed ops – **spin-identification**
- Mixing angles
- Using distillation, high stats, large operator basis, GPUs

Summary and Outlook

Summary

- **Extensive** isoscalar meson spectrum
- **Exotics** and non-exotic **hybrids**, high spin, excited states
- Carefully constructed ops – **spin-identification**
- Mixing angles
- Using distillation, high stats, large operator basis, GPUs


Outlook

- Include **glueball** operators
- A_1^{++} (0^{++}) channel
- Multi-meson operators – **resonances** and scattering
- Lighter pion masses, larger volumes, ...

Extra Slides

Prin. Corr. M_{eff}

PR D83, 111502 (2011)



$$M_{\text{eff}}(t) = -\ln [\lambda(t + \delta t) / \lambda(t)] / \delta t$$

$$\delta t = 3$$

Mixing Angle

PR D83, 111502 (2011)

